arXiv:1606.08325v1 [math.CA] 24 Jun 2016

An identity for derivatives

Ulrich Abel

Fachbereich MND
Technische Hochschule Mittelhessen
Wilhelm-Leuschner-Straße 13, 61169 Friedberg,
Germany
Ulrich.Abel@mnd.thm.de

Abstract.

We present a generalization of a formula of higher order derivatives and give a short proof.

Mathematics Subject Classification (2010): 26A06, 26A24.
Keywords: One-variable calculus, differentiation, higher derivatives.

1 Introduction

Recently, Baran e.a. [2] published the interesting identity

$$\frac{1}{n!}\sum_{k=0}^{n}(-1)^k\binom{n}{k}g^k\left(fg^{n-k}\right)^{(n)}=f\left(g'\right)^n \tag{1}$$

$(n=0,1,2,\ldots)$, for sufficiently often differentiable functions. This identity is valid in each complex commutative algebra $\mathcal{A}$ with unity and each derivation operator $D:\mathcal{A}\to\mathcal{A}$ (a linear operator with the property $D(fg)=gD(f)+fD(g)$) with the notation $D(f)=f'$ and $D^{(k)}(f)=f^{(k)}$. A proof of identity (1) is annonounced to be given in the forthcoming paper [3]. A first version with $f=1$ was found by B. Milówka [5][6] in 2005, for polynomials g. In 2012, P. Ozorka proved it for arbitrary functions g.

Such identities play a key role in deriving Markov type inequalities. M. Baran observed that the Milówka identity implies a lower estimate for the k-th derivative of polynomials considered on planar A. Markov sets.

The purpose of this note is a short transparent proof of a generalization of identity (1). We restrict ourselves to ordinary differentiation of functions on real or complex domains. One special instance is the symmetric form

$$\sum_{k=0}^{n}(-1)^k\binom{n}{k}\left(f_1g^k\right)^{(p)}\left(f_2g^{n-k}\right)^{(n-p)}=(-1)^p\,n!f_1f_2\left(g'\right)^n, \tag{2}$$

for $p=0,\ldots,n$. Note that Eq. (1) is the case $p=0$, $f_1=1$, $f_2=f$. Identity (2) can be extended to several functions (see Theorem 1).

We mention the similarity of Eq. (2) to a different identity for derivatives

$$x\sum_{k=0}^{n}\binom{n}{k}\left(x^kf\left(x\right)\right)^{(k)}\left(x^{n-k}g\left(x\right)\right)^{(n-k)}=\left(x^{n+1}f\left(x\right)g\left(x\right)\right)^{(n)}, \tag{3}$$

which is a consequence of a generalization of the Leibniz Rule [1] giving a closed form of the sum

$$\sum_{k=0}^{n}\binom{n}{k}\left(h^kf\right)^{(k)}\left(h^{n-k}g\right)^{(n-k)}.$$

2 The identity

Our notation uses multi-indices. For $\mathbf{k}=(k_1,\ldots,k_r)\in\mathbb{Z}^r$, denote $|\mathbf{k}|=k_1+\cdots+k_r$. For positive integers n, the binomial coefficient is defined by $\binom{n}{\mathbf{k}}=\binom{n}{k_1,\ldots,k_r}:=\frac{n!}{k_1!\cdots k_r!(n-|k|)!}$. Throughout the paper $\sum_{|\mathbf{k}|=n}$ means that the summation runs over all $\mathbf{k}\in\left(\mathbb{Z}_{\geq 0}\right)^r$ satisfying $|\mathbf{k}|=n$.

The following theorem contains the most general formula.

Theorem 1 *Let $n\in\mathbb{N}_0$, $r\in\mathbb{N}$, and let f_i, g_i $(i=1,\ldots,r)$ be functions which have a derivative of order n. Suppose that $\sum_{i=1}^{r}g_i=0$. Then, for $\mathbf{s}=(s_1,\ldots,s_r)\in\left(\mathbb{Z}_{\geq 0}\right)^r$,*

$$\sum_{|\mathbf{k}|=n}\binom{n}{\mathbf{k}}\prod_{i=1}^{r}\left(f_ig_i^{k_i}\right)^{(s_i)}=\begin{cases}0 & \left(|s|<n\right),\\ n!\left(\prod_{i=1}^{r}f_i\right)\prod_{i=1}^{r}\left(g_i'\right)^{s_i} & \left(|s|=n\right).\end{cases}$$

Proof. The proof is based on the observation that, by Faà di Bruno's formula, $\left(\frac{d}{dx}\right)^s f^n\left(x\right)=0$, for $s=0,\ldots,n-1$, and $\left(\frac{d}{dx}\right)^n f^n\left(x\right)=n!\left(f'\left(x\right)\right)^n$ if $f\left(x\right)=0$. Let f_i, g_i $(i=1,\ldots,r)$ be functions which have a derivative of order n in a certain point $x\in\mathbb{R}$. Furthermore, define $\mathbf{x}=(x_1,\ldots,x_r)\in\mathbb{R}^r$ and $\mathbf{1}=(1,\ldots,1)\in\mathbb{Z}^r$. We have

$$\begin{aligned}&\sum_{|\mathbf{k}|=n}\binom{n}{\mathbf{k}}\prod_{i=1}^{r}\left(f_ig_i^{k_i}\right)^{(s_i)}\left(x\right)\\ =\;&\left(\frac{\partial^{|s|}}{\partial x_1^{s_1}\cdots\partial x_r^{s_r}}\sum_{|\mathbf{k}|=n}\binom{n}{\mathbf{k}}\prod_{i=1}^{r}\left(f_ig_i^{k_i}\right)\left(x_i\right)\right)\Bigg|_{\mathbf{x}=\mathbf{1}x}\end{aligned}$$

$$= \left(\frac{\partial^{|s|}}{\partial x_1^{s_1} \cdots \partial x_r^{s_r}} \left[\left(\prod_{i=1}^{r} f_i(x_i) \right) \left(\sum_{i=1}^{r} g_i(x_i) \right)^n \right] \right) \Big|_{\mathbf{x}=\mathbf{1}x},$$

where we applied the binomial theorem. Note that $\sum_{i=1}^{r} g_i(x) = 0$ implies that

$$\frac{\partial^{|s|}}{\partial x_1^{s_1} \cdots \partial x_r^{s_r}} \left(\sum_{i=1}^{r} g_i(x_i) \right)^n \Big|_{\mathbf{x}=\mathbf{1}x} = \begin{cases} 0 & (|s| < n), \\ n! \prod_{i=1}^{r} (g_i'(x))^{s_i} & (|s| = n). \end{cases}$$

Now the desired formula follows. ■

Now we consider the particular case that the functions g_i differ only by constant factors, i.e., $g_i = c_i g$ with a function g and $c_i \in \mathbb{R}$ $(i = 1, \ldots, r)$.

Corollary 2 *Let $n \in \mathbb{N}_0$, $r \in \mathbb{N}$, and let f_i $(i = 1, \ldots, r)$ and g be functions which have a derivative of order n. Suppose that $\mathbf{c} = (c_1, \ldots, c_r) \in \mathbb{R}^r$ satisfies $|\mathbf{c}| = 0$. Let $\mathbf{s} = (s_1, \ldots, s_r) \in (\mathbb{Z}_{\geq 0})^r$. Then*

$$\sum_{|\mathbf{k}|=n} \binom{n}{\mathbf{k}} \prod_{i=1}^{r} c_i^{k_i} \left(f_i g^{k_i} \right)^{(s_i)} = \begin{cases} 0 & (|\mathbf{s}| < n), \\ n! \left(\prod_{i=1}^{r} c_i^{s_i} \right) \left(\prod_{i=1}^{r} f_i \right) (g')^{|s|} & (|\mathbf{s}| = n). \end{cases}$$

Remark 3 *In the special case $r = 2$, with $c_1 = -1 = -c_2$, we have*

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} \left(f_1 g^k \right)^{(s_1)} \left(f_2 g^{n-k} \right)^{(s_2)} = \begin{cases} 0 & (s_1 + s_2 < n), \\ (-1)^{s_1} n! f_1 f_2 (g')^n & (s_1 + s_2 = n). \end{cases}$$

We close with some direct consequences.

Example 4 *Put $f_i(x) = x^{\alpha_i}$, $g_i(x) = x^{\beta}$ $(i = 1, \ldots, r)$. Corollary 2 yields, for $\mathbf{s} = (s_1, \ldots, s_r) \in (\mathbb{Z}_{\geq 0})^r$ with $|s| = n$ and $\mathbf{c} = (c_1, \ldots, c_r) \in \mathbb{R}^r$ with $|\mathbf{c}| = 0$, the identity*

$$\sum_{|\mathbf{k}|=n} \binom{n}{\mathbf{k}} \prod_{i=1}^{r} \left(c_i^{k_i} \binom{\alpha_i + k_i \beta}{s_i} \right) = \binom{n}{\mathbf{s}} \beta^n \prod_{i=1}^{r} c_i^{s_i}. \tag{4}$$

Identity (4) can be found in Gould's collection of binomial identities [4, (6.41)]. In the special case $r = 2$ we obtain, for $s = 0, \ldots, n$,

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} \binom{\alpha_1 + k\beta}{s} \binom{\alpha_2 + (n-k)\beta}{n-s} = (-1)^s \binom{n}{s} \beta^n. \tag{5}$$

Formula (5) is an easy consequence of the fact that $\sum_{k=0}^{n} (-1)^{n-k} \binom{n}{k} k^i = 0$ $(i = 0, \ldots, n-1)$, $\sum_{k=0}^{n} (-1)^{n-k} \binom{n}{k} k^n = n!$ and $\binom{\alpha_1 + k\beta}{s} \binom{\alpha_2 + (n-k)\beta}{n-s}$ is a polynomial in the variable k of degree n with leading term $(-1)^{n-s} \beta^n k^n / (s!\,(n-s)!)$.

Example 5 *If* $f_i(x) = e^{\alpha_i x}$, $g_i(x) = e^{\beta x}$ $(i = 1, \ldots, r)$ *we obtain:*

$$\sum_{|\mathbf{k}|=n} \binom{n}{\mathbf{k}} \prod_{i=1}^{r} \left(c_i^{k_i} (\alpha_i + k_i\beta)^{s_i}\right) = \binom{n}{\mathbf{s}} \beta^n \prod_{i=1}^{r} c_i^{s_i}. \tag{6}$$

In the special case $r = 2$ *we obtain, for* $s = 0, \ldots, n$,

$$\sum_{k=0}^{n} (-1)^k \binom{n}{k} (\alpha_1 + k\beta)^s (\alpha_2 + (n-k)\beta)^{n-s} = (-1)^s n! \beta^n. \tag{7}$$

As Eq. (5) *in Ex.* 4, *formula* (7) *is obvious because* $(\alpha_1 + k\beta)^s (\alpha_2 + (n-k)\beta)^{n-s}$ *is a polynomial in the variable* k *of degree* n *with leading term* $(-1)^{n-s} \beta^n k^n$.

References

[1] U. Abel, *A generalization of the Leibniz Rule*, Amer. Math. Monthly **120:10** (2013), 924–928.
DOI 10.4169/amer.math.monthly.120.10.924

[2] Mirosław Baran, Agnieszka Kowalska, Beata Milówka and Paweł Ozorka, Identities for a derivation operator and their applications, Dolomites Research Notes on Approximation **8** (2015), 102–110.
DOI 10.14658/pupj-drna-2015-Special_Issue-10

[3] M. Baran and P. Ozorka, Derivation operators and Markov's type property for k-th iterates, 2015, in preparation.

[4] Henry W. Gould, Combinatorial Identities, Morgantown Print & Bind., Morgantown, WV (1972).

[5] B. Milówka, Markov's inequality and a generalized Plésniak condition, East Jour. of Approx. **11** (2005), 291–300.

[6] B. Milówka, Markov's property for derivatives of order k, PhD thesis (in Polish), 1–45, 2006.